

## INDEX

	PAGE
ADAMS, C. R. On the linear ordinary $q$ -difference equation . . . . .	195
ADAMS, C. R. Note on the existence of analytic solutions of non-homogeneous linear $q$ -difference equations, ordinary and partial. . . . .	626
ALBERT, A. A. On the structure of normal division algebras . . . . .	322
ALBERT, A. A. Normal division algebras in $4p^2$ units, $p$ an odd prime .	583
ALBERT, A. A. The structure of any algebra which is a direct product of rational generalized quaternion division algebras . . . . .	621
ALEXANDROFF, P. Untersuchungen über Gestalt und Lage abgeschlossener Mengen beliebiger Dimension . . . . .	101
ALTSHILLER-COURT, N. On five mutually orthogonal spheres . . . . .	613
ARCHIBALD, R. G. The impossibility of a separation of types of linear odd divisors of binary quadratic forms . . . . .	12
ARWIN, A. On cubic fields . . . . .	1
BELL, E. T. An interpretation of certain decomposable algebraic forms as functions of divisors . . . . .	429
BROWNE, E. T. On the signature of a quadratic form . . . . .	517
BURINGTON, R. S., and H. K. HOLT. Canonical forms of plane cubic curves under euclidean transformations . . . . .	52
CRAIG, C. C. The frequency function of $y/x$ . . . . .	471
EISENHART, L. P. Contact transformations . . . . .	211
EISENHART, L. P. Dynamical trajectories and geodesics . . . . .	591
GARVER, R. Quartic equations with certain groups . . . . .	47
GARVER, R. On the Brioschi normal quintic . . . . .	607
HICKEY, D. M. The equilibrium point of Green's function for an annular region	373
HILLE, E. Note on the preceding paper by Mr. Peek . . . . .	270
HOLLICROFT, T. R. Multiple lines with fixed coincident tangent planes .	451
HOLT, H. K. and R. S. BURINGTON. Canonical forms of plane cubic curves under euclidean transformations . . . . .	52
JEFFERY, R. L. The sequence of functions which define a definite integral containing a parameter. . . . .	311
KNEBELMAN, M. S. Tensors with invariant components . . . . .	339
LANE, E. P. On the fundamental transformation of surface . . . . .	457
LATIMER, C. G. On forms which repeat under multiplication . . . . .	61
LEHMER, D. H. On the multiple solutions of the Pell equation . . . . .	66
McFARLAN, L. H. A parametric problem of the calculus of variations and its treatment as a problem of Lagrange . . . . .	23
MACKIE, E. L. The Jacobi condition for a problem of Mayer with variable end-points . . . . .	393
MITRA, S. C. On certain hitherto unsolved cases of the complex multiplication of elliptic functions . . . . .	73

	PAGE
MOORE, T. W. Extended results in elimination . . . . .	92
MOORE, W. L. On the geometry of the Weddle surface . . . . .	492
PEEK, R. L. Solution to a problem in diffusion employing a non-orthogonal sine series . . . . .	265
PIEDVACHE, R. Sur la forme quadratique fondamentale d'un espace de Riemann en coordonnées normales . . . . .	272
PIEDVACHE, R. Sur le développement de Clebsch-Gordan . . . . .	281
PIERPONT, J. On the complex roots of a transcendental equation occurring in the electron theory . . . . .	81
PORITSKY, H. On certain oscillation theorems . . . . .	441
SHEFFER, I. M. Systems of linear differential equations of infinite order, with constant coefficients . . . . .	250
SHEFFER, I. M. Linear differential equations of infinite order, with poly- nomial coefficients of degree one . . . . .	345
SLOTNICK, M. M. Semi-parallel transformations of lines of curvature . . . . .	188
SRIVASTAVA, P. L. On a class of Taylor's series . . . . .	39
SRIVASTAVA, P. L. On a class of integral functions . . . . .	384
THOMAS, J. M. Riquier's existence theorems . . . . .	285
TRJITZINSKY, W. J. On quasi-analytic functions . . . . .	526
VANDIVER, H. S. On a theorem of Kummer's concerning power characters of units in a cyclotomic field . . . . .	487
VANDIVER, H. S. On the first case of Fermat's last theorem . . . . .	552
VANDIVER, H. S. An algorithm for transforming Kummer criteria in con- nection with Fermat's last theorem . . . . .	559
WARD, M. Certain expansions involving doubly infinite series . . . . .	578
WEYL, H. Der Zusammenhang zwischen der symmetrischen und der linearen Gruppe . . . . .	499
WHYBURN, W. M. On the fundamental existence theorems for differential systems . . . . .	31
WILLIAMS, K. P. A generalization of the Cauchy-Riemann equations . . . . .	206
WILSON, N. R. On finding ideals . . . . .	411
WONG, B. C. A certain point-to-line transformation in space of four dimensions . . . . .	547

---

# ERRATA

VOLUME 29, PAGE 76. Omit the Corollary.

## ON CUBIC FIELDS\*.

By A. ARWIN.

I have in this short sketch set myself the task of investigating two problems. The first of these concerns the "außerwesentlichen" primefactors for any given cubic and also higher irrationality. They are, as is known†, characterized by the fact that with a given equation  $F(x) = 0$

$$(1) \quad F(x) \equiv \prod (x) + p \cdot M(x)$$

where  $\prod (x)$  is the product of all irreducible factors of  $F(x) \pmod{p}$  and for some rational  $b$

$$\prod(b) \equiv 0, \quad \prod'(b) \equiv 0, \quad M(b) \equiv 0 \pmod{p}.$$

The general method for determining the ideal primefactors fails for such primes. We shall see how, by modification, it can be used in this case also. The other task is to find any systematic way to decide when two ideals are equivalent, and to construct units. This is done by the construction of chains. It will be seen that each equivalence from the chain gives rise to an equivalence of ideals. The converse theorem is not necessarily true. It is indeed connected with some difficulty to give a method of forming periodic developments with higher irrationalities than quadratic, and therefore it will be necessary to leave the domain  $K(1)$ , where only the quadratic irrationalities are periodic. It is however possible to form a continued set of inequalities of approximation in three dimensions and construct chains with coefficients from the cubic field  $K(\omega)$  itself, which really are periodic or have, as we shall say, periodic convergence.

As to the former problem we start from the fundamental equation

$$(2) \quad F(x) = x^3 + A_1 x^2 + A_2 x + A_3 = 0,$$

and have to treat the cases when  $F(x) \equiv 0 \pmod{p}$  has a double or a triple root. In the first case

$$F(x) \equiv (x-a)(x-b)^2 + p^i \cdot M(x),$$

\* Received March 19, 1928.

† Bachmann, P.: Allgemeine Arithmetik der Zahlkörper, p. 277.